

+EMC+ Emergency Medical Center

Your private ER

EMCemergency.com, 8245 Precinct Line Rd, # 100, N. Richland Hills, TX 76180, 817-503-8800

What Should I Do When My Child Has a Fever

What Is Fever?

Fever is a symptom--a sign that your child's body may be fighting off an infection. Most fevers are caused by common things like colds, ear infections, and bronchitis. Fever is the body's way of fighting infections. If you think your child has a fever, you should take their temperature. Temperature readings are different depending on what part of the body you use (rectum, ear, mouth). Your child has a fever if their temperature is above:

Rectal 100.4° F (38.0° C)

Oral (by mouth) 99.5° F (37.5° C)

Axillary (armpit) 98.6° F (37.0° C)

Tympanic (ear) 100.0° F (37.8° C)

Why Do We Treat a Fever?

Fevers are usually not dangerous for most children and infants over the age of 3 months. Most fevers do not lead to brain damage or death. Although some children have seizures when they have a fever, medicines used to treat fever do not always prevent these seizures. The main reason we treat fevers is to keep your child comfortable so they will eat, drink, or sleep. If your child has a mild fever but is playing, drinking fluids, and generally acting well, there is no reason to treat the fever. Medicines used to treat fever do not make the fever or infection go away faster. Medicines used to treat fever may not make your child's temperature normal. However, you should call your doctor if your child has a fever and is less than 3 months old, if the fever has lasted more than 24 hours, or if your child is also vomiting.

What Medicines Are Used to Treat Fever?

The most commonly used medicines to treat fever are acetaminophen (*Tylenol* and others) and ibuprofen (*Motrin, Advil*, others). When used as recommended, acetaminophen and ibuprofen have few side effects and are quite safe. These medicines come in drops for infants, liquid ("elixir") for toddlers, and chewable tablets for older children. The infant drops are more concentrated than the liquid elixir for toddlers. Do not switch back and forth between different products or you may give your child too much or too little medicine. Always closely read the directions on the label. DO NOT give your child aspirin for their fever as aspirin can cause Reye's syndrome.

- To avoid making mistakes, read the label before you open the bottle and again after you measure a dose.
- Do not give more or less medicine and do not give it more frequently than recommended.
- Many allergy, cold, and flu medications also contain acetaminophen or ibuprofen. Check with your pharmacist before combining medications.
- When giving your child a liquid medication, do not use standard tableware tablespoons and teaspoons because they usually are not accurate. Instead, use a measuring device such as a syringe, dropper, dosing spoon, or medicine cup.

Other Ways to Keep Your Child Comfortable

There are other ways to keep your child comfortable. These include:

- If shivering, keep your child warm until the shivering stops. If not shivering, you can remove your child's warm clothes and encourage them to drink plenty of fluids.
- Keep your child rested, quiet, and comfortable in a cool room.
- Place a cool washcloth on your child's forehead or sponge them with lukewarm water. If sponge bathing,

make sure the water doesn't get cold, and stop if your child starts to shiver.

-Never use rubbing alcohol to cool your child's skin. It can be absorbed through the skin and harm your child.

For additional information on this and other health issues, visit: MedlinePlus.gov